

QUASI-HOMOGENEOUS DOMAINS AND CONVEX AFFINE MANIFOLDS

KYEONGHEE JO

ABSTRACT. In this article, we study convex affine domains which can cover a compact affine manifold.

For this purpose, we first show that every strictly convex quasi-homogeneous projective domain has at least C^1 boundary and it is an ellipsoid if its boundary is twice differentiable. And then we show that an n -dimensional paraboloid is the only strictly convex quasi-homogeneous affine domain in $\mathbb{R}^n$ up to affine equivalence. Furthermore we prove that if a strictly convex quasi-homogeneous projective domain is C^α on an open subset of its boundary, then it is C^α everywhere.

Using this fact and the properties of asymptotic cones we find all possible shapes for developing images of compact convex affine manifolds with dimension ≤ 4 .

1. INTRODUCTION

A convex affine n -manifold is a quotient of a convex affine domain Ω in $\mathbb{R}^n$ by a discrete subgroup Γ of $\text{Aff}(n, \mathbb{R})$ acting on Ω properly discontinuously and freely. We say that a convex affine manifold is complete if $\Omega = \mathbb{R}^n$ and radiant if Ω is a cone. Particularly if M is a compact convex affine manifold then Ω becomes a divisible affine domain, i.e., its automorphism group contains a cocompact discrete subgroup acting properly. So the study of compact convex affine manifolds is equivalent to that of divisible convex affine domains and their automorphism groups.

Compact convex affine 3-manifolds were classified completely: The complete case was treated by D. Fried and W. Goldman in [16] and the radiant case was treated by T. Barbot in [1]. The remaining cases were done by H. Cho in [11], where she classified them by finding all 3-dimensional divisible convex affine domains which are not cones independently.

Recently, Y. Benoist proved in [5] that for any properly convex cone $\Omega \subset \mathbb{R}^n$ and a discrete subgroup Γ of $\text{Aff}(n, \mathbb{R})$ which divides Ω , one of the following three cases holds true:

- (i) Ω is a product,
- (ii) Ω is a symmetric cone,
- (iii) Γ is Zariski dense in $\text{GL}(n, \mathbb{R})$.

One of the oldest results in this subject is that any divisible convex affine domain Ω must be a cone if it does not contain any complete line, which was proved by

2000 *Mathematics Subject Classification.* 52A20, 57M50.

Key words and phrases. quasi-homogeneous, divisible, strictly convex, projective manifold, affine manifold.

This work was supported by Korea Research Foundation Grant (KRF-2002-070-C00010).

Vey in [42]. If Ω has a complete line it is isomorphic to $\mathbb{R}^k \times \Omega'$ for some $(n - k)$ -dimensional convex affine domain Ω' which does not contain any complete line. We see that the action of Γ on Ω induces a quotient affine action $\tilde{\Gamma}$ on Ω' . In general, $\tilde{\Gamma}$ may not be discrete any more. But Ω' still has a compact quotient by the action of $\tilde{\Gamma}$ and thus becomes just a quasi-homogeneous convex affine domain, i.e., there is a compact subset K of Ω' such that $\tilde{\Gamma}K = \Omega'$. So any divisible convex affine domain is isomorphic to the product of $\mathbb{R}^k$ and a quasi-homogeneous convex affine domain which does not contain any complete line. A good example is $\mathbb{R} \times \{(x, y) \in \mathbb{R}^2 \mid y > x^2\}$ (see [20] for details).

For this reason we study quasi-homogeneous convex affine domains to understand compact convex affine manifolds. In this article we first show that an n -dimensional paraboloid $\{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n \mid x_n > x_1^2 + x_2^2 + \dots + x_{n-1}^2\}$ is the only strictly convex quasi-homogeneous affine domain in $\mathbb{R}^n$ up to affine equivalence, in contrast with the fact that there are infinitely many quasi-homogeneous strictly convex projective domains (see [17, 28, 43]). To prove this we investigate strictly convex quasi-homogeneous projective domains, since strictly convex quasi-homogeneous affine domains are a special class of them. In general, any affine domain of $\mathbb{R}^n$ can be considered as a projective domain via the well-known equivariant embedding from $(\mathbb{R}^n, \text{Aff}(n, \mathbb{R}))$ into $(\mathbb{RP}^n, \text{PGL}(n+1, \mathbb{R}))$. So we often look at affine domains in the projective space. This enables us to use Benzécri's technique developed in convex projective domain theory [7].

Actually in section 5 we show that if Ω is a strictly convex quasi-homogeneous projective domain, then

- (i) $\partial\Omega$ is at least C^1 ,
- (ii) Ω is an ellipsoid if $\partial\Omega$ is twice differentiable,
- (iii) if $\partial\Omega$ is C^α on an open subset of $\partial\Omega$, then $\partial\Omega$ is C^α everywhere.

This implies that either any strictly convex projective domain which covers a compact projectively flat manifold is an ellipsoid or the boundary fails to be twice differentiable on a dense subset, which is a generalization of the 2-dimensional result of Kuiper [29].

For divisible case, a similar result has been proved independently by Y. Benoist in [4] and he claimed in [5] that any quasi-homogeneous strictly convex projective domain which is not an ellipsoid, has a discrete automorphism group. We will also show the claim in section 5.3.

Another earlier result related to this subject is that if Ω is a strictly convex (in the sense that Hessian is positive definite) domain with C^3 boundary which has a cofinite volume discrete subgroup action, then Ω is an ellipsoid [13].

To conclude that any strictly convex quasi-homogeneous affine domain is affinely equivalent to a paraboloid, we show in section 5.2 that its boundary is homogeneous and thus C^∞ . (Then the domain is projectively equivalent to an ellipsoid by the above result (ii) about quasi-homogeneous strictly convex projective domain and this implies that it is affinely equivalent to a paraboloid.) To do this we first prove that every strictly convex domain has an invariant direction under the action of its affine automorphism group, that is, it contains a half line which is invariant under the action of the linear part of its automorphism group. Generally, we will show in section 4 that every convex quasi-homogeneous affine domain Ω is foliated by cosets of the *asymptotic cone* which is the maximal cone contained in Ω and invariant under the action of the linear part of its automorphism group.

Using this asymptotic cone we find all the possible quasi-homogeneous convex affine domains with dimension ≤ 3 . Then we find all the possible shapes for a universal covering space of compact convex affine manifold with dimension ≤ 4 . Finally, we give a partial result to the Markus conjecture which says that “A compact affine manifold is complete if and only if it has parallel volume.” This conjecture was proved under some additional conditions about holonomy group by J. Smillie, D. Fried, W. Goldman, M. Hirsch, Carrière and so on, see [8], [9], [14] [15], [21], [22], [30], [38]. In this article we show that any compact convex affine manifold with dimension ≤ 4 is complete if it has parallel volume.

2. PRELIMINARIES

We present here some of the basic materials from Benzécri’s convex domain theory that we will need later. We begin with some definitions.

The real projective space $\mathbb{RP}^n$ is the quotient space of $\mathbb{R}^{n+1} \setminus \{0\}$ by the action of $\mathbb{R}^* = \mathbb{R} \setminus \{0\}$. In an affine space, we usually denote the affine subspace generated by a subset A by $\langle A \rangle$. So we will use the same notation for a subset of $\mathbb{RP}^n$, i.e., for each subset B of $\mathbb{RP}^n$ $\langle B \rangle$ means the projectivization of the affine subspace generated by $\pi^{-1}(B)$ in $\mathbb{R}^{n+1}$, where π is the quotient map from $\mathbb{R}^{n+1} \setminus \{0\}$ onto $\mathbb{RP}^n$ and we will call $\langle B \rangle$ the *support* of B .


A *quasi-homogeneous* affine (respectively, projective) domain is an open subset Ω of $\mathbb{R}^n$ (respectively, $\mathbb{RP}^n$) which has a compact subset $K \subset \Omega$ and a subgroup G of $\text{Aut}(\Omega)$ such that $GK = \Omega$, where $\text{Aut}(\Omega)$ is a subgroup of $\text{Aff}(n, \mathbb{R})$ (respectively, $\text{PGL}(n+1, \mathbb{R})$) consisting of all affine (respectively, projective) transformations preserving Ω . Sometimes we say that G acts on Ω *syndetically*. It follows that both homogeneous domains and divisible domains are quasi-homogeneous.

As stated in the introduction, we can consider an affine domain as a projective domain and it is obvious that every quasi-homogeneous affine domain is a quasi-homogeneous projective domain. But a quasi-homogeneous projective domain is not a quasi-homogeneous affine domain even if it is contained in an affine patch of a projective space. For examples, any quasi-homogeneous strictly convex projective domain which is not an ellipsoid, is not a quasi-homogeneous affine domain (this will be proved in section 7) and in fact there exist infinitely many such quasi-homogeneous strictly convex projective domains. (See [17] and [43] for the 2-dimensional case and [2] for arbitrary dimensional cases.)

For an affine domain $\Omega \subset \mathbb{R}^n$, we denote the group of all affine transformations preserving Ω by $\text{Aut}_{\text{aff}}(\Omega)$ to distinguish it from the group of all projective transformations preserving Ω . Note that $\text{Aut}_{\text{aff}}(\Omega)$ is a closed subgroup of $\text{Aut}(\Omega)$. Also usually we denote the boundary of a domain Ω by $\partial\Omega$, but sometimes we will use the $\partial_a\Omega$ and $\partial_p\Omega$ when it is necessary to avoid ambiguity and call them an affine boundary and a projective boundary of Ω , respectively. Note that $\partial_a\Omega$ is a subset of $\partial_p\Omega$ and in fact $\partial_a\Omega = \mathbb{R}^n \cap \partial_p\Omega$.

For a convex projective domain $\Omega \subset \mathbb{RP}^n$, we define a relation $\sim$ on $\overline{\Omega}$ by defining $x \sim y$ if either $x = y$ or $x \neq y$ and $\overline{\Omega}$ has an open line segment l containing both x and y . It follows from the convexity of Ω that $\sim$ is an equivalence relation : its symmetry is obvious by definition. See Fig. 1 for its transitivity.

An equivalence class with respect to $\sim$ is called a *face* of Ω and we will call the closure of a face a *closed face* of Ω . Note that a face is relatively open in its support and $\overline{\Omega}$ is a disjoint union of all faces. A convex domain Ω in $\mathbb{RP}^n$ is

FIGURE 1. Transitivity of $\sim$

called *properly convex* if there is no non-constant projective map of $\mathbb{R}$ into Ω and *strictly convex* if $\partial\Omega$ contains no line segment. From these definitions we see that any strictly convex domain is a properly convex domain.

It is clear that the intersection of any family of convex sets is again convex. Therefore, for any subset A there is a smallest convex set containing A , namely, the intersection of all convex sets containing A . This convex set is called the convex set spanned by A , or the convex hull of A , and is denoted by $CH(A)$.

An affine subspace $\mathcal{S}$ is called a *supporting subspace* of a convex domain Ω in $\mathbb{RP}^n$ if following conditions are satisfied :

- (i) $\mathcal{S} \cap \overline{\Omega} \neq \emptyset$, and
- (ii) for any face F of Ω such $F \cap \mathcal{S} \neq \emptyset$, F is entirely contained in $\mathcal{S}$.

We see immediately that the support of a face is always a supporting subspace and any hyperplane H tangent to $\partial\Omega$ is a supporting subspace of Ω .

Example 2.1.

- (i) Let $\Omega = \{(x, y, z) \in \mathbb{R}^3 \mid y > x^2, z > 0\}$. Then the z -axis is the support of a face $\{(0, 0, z) \in \mathbb{R}^3 \mid z > 0\}$, the y -axis is neither a support of any face nor a supporting subspace of Ω , and the x -axis is not a support of any face but a supporting subspace of Ω .
- (ii) Let Ω be the standard simplex in $\mathbb{R}^n$, that is, its vertices are $\{0, e_1, \dots, e_n\}$. Then $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 1\}$ is a supporting subspace of Ω . (Note that this is not a quasi-homogeneous affine domain but a quasi-homogeneous projective domain.)

The following definitions are equivalent to Benzécri's original definitions in [7].

Definition 2.2.

- (i) Let Ω be a properly convex domain in $\mathbb{RP}^n$ and let Ω_1 and Ω_2 be convex domains in $\langle\Omega_1\rangle$ and $\langle\Omega_2\rangle$ respectively. Ω is called a *convex sum* of Ω_1 and Ω_2 , which will be denoted by $\Omega = \Omega_1 + \Omega_2$, if $\langle\Omega_1\rangle \cap \langle\Omega_2\rangle = \emptyset$ and Ω is the union of all open line segments joining points in Ω_1 to points in Ω_2 .
- (ii) A k -dimensional face F of an n -dimensional convex domain Ω is called *conic* if there exist $n - k$ supporting hyperplanes $H_1, H_2, \dots, H_{n-k}$ such that

$$H_1 \supsetneq H_1 \cap H_2 \supsetneq \dots \supsetneq H_1 \cap \dots \cap H_{n-k} = \langle F \rangle.$$

- (iii) Let Ω be a properly convex domain in $\mathbb{RP}^n$. We say that Ω has an *osculating ellipsoid* at $p \in \partial\Omega$ if there exist a suitable affine chart and a basis such that the local boundary equation on some neighborhood of $p = (0, \dots, 0)$

is expressed by $x_n = f(x_1, \dots, x_{n-1})$ and

$$\lim_{(x_1, \dots, x_{n-1}) \rightarrow 0} \frac{f(x_1, \dots, x_{n-1})}{x_1^2 + \dots + x_{n-1}^2} = 1.$$

Remark 2.3. From Definition 2.2 (ii), any $(n-1)$ -dimensional face is conic.

Example 2.4.

- (i) Let Ω be a simplex. Then every face of Ω is conic.
- (ii) Let $\Omega = \{(x, y, z) \in \mathbb{R}^3 \mid y > x^2, z > 0\}$. Then it is a quasi-homogeneous affine (and so projective) domain and the subset $\{(0, 0, z) \in \mathbb{R}^3 \mid z > 0\}$ is a non-conic face of Ω .

2.1. Spaces of convex bodies in $\mathbb{RP}^n$. In 1960, Benzécri developed projective convex body theory in [7]. We recall here some important results among them, which can be also found in [19]. We will follow Goldman's notation [19].

First, we call a subset $K \subset \mathbb{RP}^n$ a *convex body* if K is the closure of a properly convex domain of $\mathbb{RP}^n$. Equivalently, K is a convex body if K is a closed convex subset with nonempty interior of $\mathbb{RP}^n$ which does not contain any complete line.

Let $\mathcal{C}(n)$ denote the set of all convex bodies in $\mathbb{RP}^n$, with the topology induced from the Hausdorff metric on the set of all closed subsets of $\mathbb{RP}^n$. Let

$$\mathcal{C}_*(n) = \{(K, x) \in \mathcal{C}(n) \times \mathbb{RP}^n \mid x \in \text{int}(K)\}$$

be the corresponding set of pointed convex bodies with topology induced from the product topology on $\mathcal{C}(n) \times \mathbb{RP}^n$.

Theorem 2.5 (Benzécri). *$PGL(n+1, \mathbb{R})$ acts properly and syndetically on $\mathcal{C}_*(n)$. In particular the quotient $\mathcal{C}_*(n)/PGL(n+1, \mathbb{R})$ is a compact Hausdorff space.*

Remark 2.6. While the quotient $\mathcal{C}_*(n)/PGL(n+1, \mathbb{R})$ is Hausdorff, the space of equivalence classes of convex bodies $\mathcal{C}(n)/PGL(n+1, \mathbb{R})$ is not Hausdorff. Some basic examples can be found in [7] and in C.16 of [19].

Using the above theorem, he proved the following important result.

Theorem 2.7 (Benzécri). *Let $\Omega \subset \mathbb{RP}^n$ is a quasi-homogeneous properly convex domain. Then the corresponding point $[\bar{\Omega}] \in \mathcal{C}(n)/PGL(n+1, \mathbb{R})$ is closed, that is, if $g_n \Omega$ converges to some properly convex domain $\Omega' \subset \mathbb{RP}^n$ for some $\{g_n\} \subset PGL(n+1, \mathbb{R})$, then Ω is projectively equivalent to Ω' .*

Theorem 2.8 (Benzécri). (i) *Let Ω be a properly convex projective domain in $\mathbb{RP}^n$ and F be a conic face of Ω . Then there exist a projective subspace L of $\mathbb{RP}^n$ and projective automorphisms $\{h_i\}$ of $\mathbb{RP}^n$ such that $\{h_i \Omega\}$ converges to $F + B$ for some properly convex domain B in L .*

(ii) *If a properly convex domain Ω has an osculating ellipsoid Q , then there exists a sequence $\{g_n\} \subset PGL(n+1, \mathbb{R})$ such that $g_n \Omega$ converges to Q .*

(iii) *Let $\Omega = \Omega_1 + \Omega_2$. Then Ω is quasi-homogeneous (respectively, homogeneous) if and only if Ω_i is quasi-homogeneous (respectively, homogeneous) for each i .*

(iv) *Let Ω be a quasi-homogeneous properly convex domain in $\mathbb{RP}^n$ and L a linear subspace of $\mathbb{RP}^n$ of dimension r such that $L \cap \Omega$ has a conic face F . Then there exists a section which is projectively equivalent to an r -dimensional properly convex domain $F+B$ for some suitable properly convex domain B of dimension $r - (\dim(F) + 1)$.*

Here a section means an intersection of a projective subspace and Ω .

Corollary 2.9. *Any quasi-homogeneous properly convex projective domain which is not strictly convex has a triangle as a 2-dimensional section.*

Proof. Let Ω be a domain satisfying the hypothesis. Then $\partial\Omega$ contains a line segment l . Considering a 2-dimensional plane P which intersects Ω and contains l , we see that l is a conic face of $P \cap \Omega$. By Theorem 2.8 (iv), we get the desired result. $\square$

3. $\text{Aut}(\Omega)$ AND ITS LIMIT PROJECTIVE TRANSFORMATIONS

Any properly convex projective domain is projectively equivalent to a bounded convex domain in an affine space. So we can define on any properly convex projective domain a complete continuous metric which is invariant under the action of $\text{Aut}(\Omega)$. This metric is called the Hilbert metric and defined as follows.

Definition 3.1. Let Ω be a bounded convex domain in $\mathbb{R}^n$. For any two different points $p_1, p_2 \in \Omega$, we define $d_H(p_1, p_2)$ to be the logarithm of the absolute value of the cross ratio of (s_1, s_2, p_1, p_2) , where s_1 and s_2 are the points in which the line $\overline{p_1 p_2}$ intersects $\partial\Omega$. For $p_1 = p_2$, we define $d_H(p_1, p_2) = 0$.

Since $\text{PM}(n+1, \mathbb{R})$, which is the projectivization of the group of all $(n+1)$ by $(n+1)$ matrices, is a compactification of $\text{PGL}(n+1, \mathbb{R})$, any infinite sequence of non singular projective transformations contains a convergent subsequence. Note that the limit projective transformation may be singular. For a singular projective transformation g we will denote the projectivization of the kernel and range of g by $K(g)$ and $R(g)$. Then g maps $\mathbb{RP}^n \setminus K(g)$ onto $R(g)$ and the images of any compact set in $\mathbb{RP}^n \setminus K(g)$ under the convergent sequence g_i , converge uniformly to the images under the limit transformation g of g_i (see [7] and [23]).

Lemma 3.2. *Let Ω be a properly convex domain in $\mathbb{RP}^n$ and let $\{g_i\} \subset \text{Aut}(\Omega)$ be a sequence converging to a singular projective transformation $g \in \text{PM}(n+1, \mathbb{R})$. Then,*

- (i) $K(g)$ is a supporting subspace of Ω ,
- (ii) $R(g) = \langle R(g) \cap \partial\Omega \rangle$ and $R(g)$ is a supporting subspace of Ω .

Proof. In [7, p310–311], Benzécri proved that $K(g)$ is a supporting subspace and $R(g) = \langle R(g) \cap \partial\Omega \rangle$. So it suffices to show that $R(g)$ is a supporting subspace of Ω . Suppose not. We can choose two points $\{x, y\}$ which are contained in $\partial\Omega \cap R(g)^c$ such that $\overline{xy} \cap R(g)$ is a point, that is, $\overline{xy}$ meets $R(g)$ transversally. Then we can find a small neighborhood U of $R(g) \cap \partial\Omega$ in $\mathbb{RP}^n$ which does not contain any ε -ball with respect to the Hilbert metric on Ω . But any compact ε -ball neighborhood $B(z, \varepsilon)$ of a point z in Ω is mapped into $U \cap \overline{\Omega}$ under the g_i action for sufficiently large i , since $g_i(B(z, \varepsilon))$ converges uniformly to $g(B(z, \varepsilon)) \subset R(g) \cap \partial\Omega$. This contradiction completes the proof. $\square$

Remark 3.3.

- (1) Note that neither $K(g)$ nor $R(g)$ intersects Ω since $\dim K(g) < n$ and $\dim R(g) < n$ in this situation.
- (2) It follows from (ii) of the above lemma that $R(g) \cap \partial\Omega$ is a closed face of Ω . In fact, we can show that any supporting subspace L satisfying $L = \langle L \cap \partial\Omega \rangle$ is always a support of a face.

Lemma 3.4. *Let Ω be a quasi-homogeneous properly convex domain in $\mathbb{RP}^n$ and G a subgroup of $\text{Aut}(\Omega)$ acting on Ω syndetically. Then we get the following :*

- (i) *For each point $p \in \partial\Omega$, there exists a sequence $\{g_i\} \subset G$ and $x \in \Omega$ such that $g_i(x)$ converges to p .*
- (ii) *For any accumulation point g of $\{g_i\}$, $R(g)$ is the projective subspace generated by the face containing p .*

Proof.

- (i) Choose a straight line segment ℓ in $\overline{\Omega}$ with initial point in Ω and endpoint p . Let K be a compact generating domain of Ω . Then ℓ intersects an infinite series of different images of K ; $g_1(K), g_2(K), \dots$. Choose $x_i \in \ell \cap g_i(K)$ for all i . Then we have $g_i^{-1}(x_i) \in K$ and thus we can choose $x \in K$ which is one of the accumulation points of $\{g_i^{-1}(x_i)\}$ since K is compact. We may assume $\{g_i\}$ converges to g . Since $d_H(g_i(x), x_i) = d_H(x, g_i^{-1}(x_i))$ converges to 0, $d(g_i(x), x_i)$ also converges to 0, where d_H is a Hilbert metric and d is the standard metric of $\mathbb{RP}^n$. We see that $d(g_i(x), p) \leq d(g_i(x), x_i) + d(x_i, p)$ and this implies that $g_i(x)$ converges to p .
- (ii) Obviously g is a singular projective transformation. Let F be the face of Ω which contains p . Then by Lemma 3.2 $R(g)$ contains F and $R(g) = \langle R(g) \cap \partial\Omega \rangle$. Since p must be an interior point of $R(g) \cap \partial\Omega$, we get $R(g) = \langle F \rangle$.

□

Lemma 3.5. *Let $\{f_i\}$ be a sequence in $\text{Aff}(n, \mathbb{R}) (< PGL(n+1, \mathbb{R}))$. Suppose that f_i converges to $f \in PM(n+1, \mathbb{R})$ with $R(f) \cap \mathbb{R}^n \neq \emptyset$. Then $K(f) \cap \mathbb{R}^n = \emptyset$.*

Proof. Suppose $K(f) \cap \mathbb{R}^n \neq \emptyset$ and choose $x \in K(f) \cap \mathbb{R}^n$. For any one-dimensional affine subspace L such that $L \cap K(f) = \{x\}$, $L \setminus \{x\}$ is mapped into a point in $R(f)$ by f . Since every one-dimensional affine subspace through x meets $\partial\mathbb{R}^n = \mathbb{RP}_\infty^{n-1}$, for any $p \in R(f)$ there exist $q \in \partial\mathbb{R}^n \setminus K(f)$ such that $f_i(q)$ converges to $p = f(q)$. Now we choose any point y in $\mathbb{R}^n \setminus K(f)$ such that $f(y) \in R(f) \cap \mathbb{R}^n$. Then the one-dimensional affine subspace l_{xy} containing x and y intersects $\partial\mathbb{R}^n$ at a point z . Since $l_{xy} \cap K(f) = \{x\}$, z is not contained in $K(f)$ and so $f_i(z)$ converges to $f(z) = f(y)$. But this is impossible since $f_i(z) \in f_i(\partial\mathbb{R}^n) = \partial\mathbb{R}^n$ for all i and $f(y) \in \mathbb{R}^n$. This completes the proof. □

Lemma 3.6. *Let Ω be a quasi-homogeneous properly convex affine domain in $\mathbb{R}^n$. Then $\partial_a\Omega$ does not have any bounded face with dimension $k > 0$.*

Proof. Let L be a supporting subspace of $\dim k$ such that $\mathcal{S} = \partial_a\Omega \cap L$ is a bounded closed subset of L with nonempty interior. By Lemma 3.4, there exists a sequence $\{f_i\} \subset \text{Aut}_{\text{aff}}(\Omega)$ which converges to $f \in PM(n+1, \mathbb{R})$ with $R(f) = L$ since $\text{int}\mathcal{S}$ is a face of Ω . Then Lemma 3.5 implies that $K(f) \subset \partial\mathbb{R}^n$ and $K(f) \neq \partial\mathbb{R}^n$ since $k > 0$. Note that $f_i(\partial\mathbb{R}^n) = \partial\mathbb{R}^n$ for each i and this implies $f(\partial\mathbb{R}^n \setminus K(f)) \subset \partial\mathbb{R}^n$.

Let K be the linear subspace of all points with direction in $K(f)$. Then K is $(n-k)$ -dimensional. Choose any k -dimensional linear subspace N which is transversal to K and consider a projection p from $\mathbb{R}^n$ to N satisfying $p(x+k) = x$ for any $x \in \mathbb{R}^n$ and any $k \in K$. Suppose $p(\Omega)$ is not bounded in N . Then there exist sequences $\{x_n\} \subset N$ and $\{y_n\} \subset \Omega$ such that $y_n \in x_n + K$ and x_n converges to a point x_∞ in $\overline{N} \cap \partial\mathbb{R}^n \subset \partial\mathbb{R}^n \setminus K(f)$, where $\overline{N}$ means the closure of N in $\mathbb{RP}^n$. Observe that $f(x_n) = f(y_n)$ since y_n lies in the projection of $\pi^{-1}(\langle x_n, K(f) \rangle)$. But

$f(x_n)$ converges to a point $f(x_\infty)$ in $\overline{L} \cap \partial \mathbb{R}^n$ and $f(y_n)$ is a point $\mathcal{S} = L \cap \partial \Omega$. This is a contradiction. So $p(\Omega)$ must be bounded in N . This implies that Ω must lie between two parallel hyperplanes. But it is known that any quasi-homogeneous affine domain can not lie between parallel hyperplanes (see [32]). $\square$

For a properly convex open cone Vey showed in [42] that $\text{Aut}_{\text{aff}}(\Omega)$ acts properly on Ω using the properties of the characteristic function of Ω . In fact we can prove the following.

Proposition 3.7. *Let Ω be a properly convex domain in $\mathbb{RP}^n$ and G be a closed subgroup of $\text{Aut}(\Omega)$. Then G acts properly on Ω .*

Proof. Let K and L be compact subsets of Ω . Define $C(K, L) \subset \text{Aut}(\Omega)$ by

$$C(K, L) = \{g \in G \mid gK \cap L \neq \emptyset\}.$$

Let $\{g_n\} \subset C(K, L)$ be a sequence converging to $g \in \text{PM}(n+1, \mathbb{R})$. By the definition of $C(K, L)$, there exists $x_n \in K$ such that $g_n x_n \in L$ for each n . Since K and L are compact, we may assume that x_n converges to $x \in K$ and $g_n x_n$ converges to $y \in L$ by choosing a subsequence if necessary. Then,

$$\begin{aligned} d_H(g_n x, y) &\leq d_H(g_n x, g_n x_n) + d_H(g_n x_n, y) \\ &= d_H(x, x_n) + d_H(g_n x_n, y), \end{aligned}$$

$$\lim_{n \rightarrow \infty} d_H(g_n x, y) = 0, \text{ i.e. } \lim_{n \rightarrow \infty} g_n x = y.$$

Suppose g is singular. By Lemma 3.2 (i), $x \notin K(g)$ and thus $gx = y$ is contained in $R(g)$. This contradicts Lemma 3.2 (ii). Therefore g is nonsingular and so $g \in C(K, L)$. This implies that $\overline{C(K, L)} = C(K, L)$ in $\text{PM}(n+1, \mathbb{R})$ and we conclude that $C(K, L)$ is compact. $\square$

The following lemma is actually an immediate corollary of the above proposition, but we give a direct proof here.

Lemma 3.8. *Let Ω be a properly convex projective domain in $\mathbb{RP}^n$ and G be a closed subgroup of $\text{Aut}(\Omega) < \text{PGL}(n+1, \mathbb{R})$. Then each G -orbit is a closed subset of Ω .*

Proof. Let $x \in \Omega$ and $y \in \overline{Gx} \cap \Omega$. Choose a sequence $\{g_n\}$ of automorphisms such that $\lim_{n \rightarrow \infty} g_n x = y$. Then there exists a subsequence $\{g_{n_k}\}$ of $\{g_n\}$ converging to some $g \in \text{PM}(n+1, \mathbb{R})$ by compactness of $\text{PM}(n+1, \mathbb{R})$. Suppose g is singular. Then by Lemma 3.2 we know $x \notin K(g)$ and this implies $y \in R(g)$. This contradicts Lemma 3.2. So g can not be singular, that is, $g \in \text{Aut}(\Omega)$. Thus $y = gx \in Gx$ since G is closed in $\text{Aut}(\Omega)$. $\square$

The following theorem is well-known and a proof is given in Theorem 6.5.1. of [33].

Theorem 3.9. *Let Γ be a group of isometries of a metric space X and d_Γ the orbit space distance function*

$$d_\Gamma : X/\Gamma \times X/\Gamma \rightarrow \mathbb{R}$$

defined by the formula

$$d_\Gamma(\Gamma x, \Gamma y) = \text{dist}(\Gamma x, \Gamma y) = \inf\{d(z_1, z_2) \mid z_1 \in \Gamma x \text{ and } z_2 \in \Gamma y\}.$$

Then d_Γ is a metric on X/Γ if and only if each Γ -orbit is a closed subset of X .

By Lemma 3.8 and Theorem 3.9, we obtain a metric d_G on the orbit space $G \backslash \Omega$ using d_H as follows:

$$(1) \quad d_G(Gx, Gy) = \inf_{g, g' \in G} d_H(gx, g'y).$$

Since d_H is invariant by projective automorphism, we have

$$d_G(Gx, Gy) = \inf_{g \in G} d_H(x, gy).$$

Up to now we have shown that the Hilbert metric on a properly convex domain Ω descends to the orbit space $G \backslash \Omega$ if G is a closed subgroup of $\text{Aut}(\Omega)$. In addition, we can show that the distance between the two orbits Gx and Gy is realized by the Hilbert distance between two points $x_0 \in Gx$ and $y_0 \in Gy$ in Ω .

Lemma 3.10. *Let G be a closed subgroup of $PGL(n+1, \mathbb{R})$. Then for each pair $(Gx, Gy) \in G \backslash \Omega$, there exists $g_{x,y} \in G$ such that $d_G(Gx, Gy) = d_H(x, g_{x,y}y)$.*

Proof. By (1), there exists a sequence $\{g_n\} \subset G$ such that $d_H(x, g_n y)$ converges to $d_G(Gx, Gy)$. For any $r > d_G(Gx, Gy)$, there exists $N > 0$ such that $g_n y \in B(x, r)$ whenever $n \geq N$. We may assume that $\{g_n\}$ converge to a projective transformation $g \in PM(n+1, \mathbb{R})$ by taking a subsequence if necessary. If g is singular then $g_n y$ converges to some point in $\partial\Omega$. This is a contradiction. Therefore $\{g_n\}$ converges to $g_{x,y} \in G$ and so we get

$$d_H(x, g_{x,y}y) = \lim_{n \rightarrow \infty} d_H(x, g_n y)$$

since $g_n y$ converges to $g_{x,y}y \in \Omega$. $\square$

4. ASYMPTOTIC CONE

As a special subclass of quasi-homogeneous convex projective domains, quasi-homogeneous convex affine domains have some important properties which distinguish them from other projective domains. They contain a cone invariant under the action of linear parts of their automorphism groups, which is called an asymptotic cone. This terminology was originally introduced by Vey in [42].

Definition 4.1. Let Ω be a convex domain in $\mathbb{R}^n$. The *asymptotic cone* of Ω is defined as follows:

$$AC(\Omega) = \{u \in \mathbb{R}^n \mid x + tu \in \Omega, \text{ for all } x \in \Omega, t \geq 0\}.$$

By the convexity of Ω , for any $x_0 \in \Omega$,

$$AC(\Omega) = AC_{x_0}(\Omega) := \{u \in \mathbb{R}^n \mid x_0 + tu \in \Omega, \text{ for all } t \geq 0\}.$$

Note that $AC(\Omega)$ is a properly convex closed cone in $\mathbb{R}^n$ if Ω is properly convex.

Remark 4.2. (i) *It is well known that there is no bounded quasi-homogeneous affine domain (see [1]). So we see that $AC(\Omega)$ is not empty if Ω is quasi-homogeneous.*
(ii) *Actually, the asymptotic cone of Ω is the maximal closed cone which can be contained in Ω .*

Now we will show that any properly convex quasi-homogeneous affine domain is foliated by cosets of its asymptotic cone. To prove this, we need the following result (see [42]).

Theorem 4.3 (Vey). *Let Ω be a properly convex open cone in $\mathbb{R}^n$ and $G < GL(n, \mathbb{R})$ acts syndetically on Ω . If L is a G -stable subspace generated by the cone $\overline{\Omega} \cap L$, the sections $\Omega_x = \Omega \cap (x + L)$ are cones for each $x \in \Omega$.*

Let L be the linear subspace of $\mathbb{R}^n$ which is generated by $AC(\Omega)$ and Ω_x the intersection of Ω and the affine subspace $x + L$, that is, $\Omega_x = \Omega \cap x + L$. Then we see immediately that $x + AC(\Omega) \subset \Omega_x$ for all $x \in \Omega$ since Ω is convex. Actually we can show that Ω_x is the translation of $AC(\Omega)$ for all $x \in \Omega$. More precisely, any section of Ω which is cut in parallel with L is itself a cone, which is exactly an asymptotic cone.

Theorem 4.4. *Let Ω be a properly convex quasi-homogeneous affine domain in $\mathbb{R}^n$. Then Ω admits a parallel foliation by cosets of the asymptotic cone of Ω , i.e., Ω_x is a translation of the asymptotic cone of Ω for all $x \in \Omega$.*

Proof. In the vector space $\mathbb{R}^{n+1} = \{(x, t) \mid x \in \mathbb{R}^n, t \in \mathbb{R}\}$, the affine hyperplane $L_1 = \{(x, t) \mid x \in \mathbb{R}^n, t = 1\}$ is isomorphic to $\mathbb{R}^n$. So we can consider Ω as an open set of L_1 , and G as a subgroup of $GL(n+1, \mathbb{R})$ conserving L_1 and L_0 by the following correspondence:

$$(i, \rho) : (\mathbb{R}^n, \text{Aff}(n, \mathbb{R})) \rightarrow (\mathbb{R}^{n+1}, GL(n+1, \mathbb{R})),$$

$$i(x_1, \dots, x_n) = (x_1, \dots, x_n, 1),$$

$$\rho(A, a) = \begin{pmatrix} A & a \\ 0 & 1 \end{pmatrix}.$$

We define a properly convex open cone Ω_1 of $\mathbb{R}^{n+1}$:

$$\Omega_1 = \{(x, t) \mid t > 0, \frac{x}{t} \in \Omega\}.$$

We easily see that $\overline{\Omega}_1 \cap L_0$ is isomorphic to the asymptotic cone of Ω . Let G_1 be the subgroup of $GL(n+1, \mathbb{R})$ generated by G and $2^m I_{n+1}$ ($m \in \mathbb{Z}$). Then G_1 acts on Ω_1 and preserve L_0 . And we have the following equalities:

$$\begin{aligned} \Omega_x &= \Omega_1 \cap (x + \langle L_0 \cap \overline{\Omega}_1 \rangle) \\ &= \Omega \cap (x + \langle L_0 \cap \overline{\Omega}_1 \rangle) \\ &= \Omega \cap (x + \langle AC(\Omega) \rangle) \quad \text{for any } x \in \Omega. \end{aligned}$$

By applying Theorem 4.3 to $L = \langle L_0 \cap \overline{\Omega}_1 \rangle$, we see Ω_x must be a cone and thus must be the asymptotic cone of Ω by convexity. $\square$

From Theorem 4.4 we get the following corollary which was originally proved by Vey [42].

Corollary 4.5. *Let Ω be a properly convex quasi-homogeneous affine domain in $\mathbb{R}^n$. If the dimension of $AC(\Omega)$ is equal to n , then Ω is a cone.*

Theorem 4.4 implies that for each $x \in \Omega$ there exists a point $s(x)$ in its boundary such that $\Omega_x = AC(\Omega) + s(x)$. So we get a continuous map $s : \Omega \rightarrow \partial\Omega$ and an one parameter group of homeomorphisms of Ω by the following formula:

$$c_t(x) = s(x) + e^t(x - s(x)) \text{ for } t \in \mathbb{R}, x \in \Omega.$$

This flow preserves every asymptotic section Ω_x and gives a nonvanishing vector field on Ω which is similar to the Euler vector field of any cone.

- Definition 4.6.** (i) We call $p \in \partial\Omega$ an *asymptotic cone point* of a properly convex affine domain Ω if p is a cone point of Ω_x for some $x \in \Omega$, that is, $p = s(x)$ for some $x \in \Omega$.
- (ii) We call $p \in \partial\Omega$ an *extreme point* of a properly convex projective domain Ω if p is not expressed by a convex sum of any two points in $\partial\Omega$, or equivalently, a zero dimensional face of Ω .

It is well-known that if Ω is a properly convex domain in $\mathbb{RP}^n$ then $\overline{\Omega}$ is the closed convex hull of the set of its extreme points (the Krein-Milman theorem; see [36]). The following proposition says that asymptotic cone points of any properly convex affine domain are the only extreme points in $\mathbb{R}^n$, that is, all other extreme points are at infinity.

Proposition 4.7. *Let Ω be a quasi-homogeneous properly convex affine domain in $\mathbb{R}^n$. Let E be the set of all extreme points of $\overline{\Omega} \subset \mathbb{RP}^n$ and S the set of all asymptotic cone points $s(x)$. Then*

$$E = S \cup E_\infty$$

where E_∞ denote the set of all extreme points of $\overline{AC(\Omega)} \cap \mathbb{RP}_\infty^{n-1} = \overline{\Omega} \cap \mathbb{RP}_\infty^{n-1}$.

Proof.

- (i) $S \subset E$

Suppose not. Then there exists an open line segment $l \subset \overline{\Omega}$ which contains a point $s(x)$ for some $x \in \Omega$. Note that $\overline{ab}$ is obviously contained in $\partial\Omega$ since $s(x) \in \partial\Omega$. Let F be the face containing $s(x)$. By Lemma 3.6, we know that F cannot be bounded and so it contains an asymptotic direction v . This means that $s(x) + tv \in F$ for all $t \geq 0$. Since F is open in $\langle F \rangle$, we can choose a negative real number t such that $s(x) + tv \in F$. But this contradicts that $s(x)$ is an asymptotic cone point. Therefore we conclude that $S \subset E$.

- (ii) $E_\infty \subset E$

This follows from the fact that any extreme point of a face of Ω is an extreme point of Ω .

- (iii) $E \subset S \cup E_\infty$

Let $e \in \mathbb{R}^n$ be an extreme point of Ω , that is, $e \in E \setminus E_\infty$. Consider the affine cone $e + AC(\Omega)$. The convexity of $\overline{\Omega}$ implies that $e + AC(\Omega)$ is a subset of $\overline{\Omega}$. If $(e + AC(\Omega)) \cap \Omega \neq \emptyset$, then $e + AC(\Omega)$ must be a section of Ω by Theorem 4.4 and thus $e \in S$. If $e + AC(\Omega) \subset \partial\Omega$, then $e + AC(\Omega)$ must be a closed face of Ω by Theorem 4.4 again. Choose a sequence $\{g_n\} \subset \text{Aut}_{\text{aff}}(\Omega)$ converging to $g \in \text{PM}(n+1, \mathbb{R}^n)$ such that $R(g) = \{e\}$. Then $K(g) = \partial\mathbb{R}^n$ by Lemma 3.5 and so $g_n(s)$ converges to e for any $s \in S$. Since $g_n(s) \in S$, we get $e \in \overline{S} \cap \mathbb{R}^n$.

Suppose that $S' = (\overline{S} \cap \mathbb{R}^n) \setminus S$ is not empty. Since $\overline{S}$, $\mathbb{R}^n$ and S are preserved by $\text{Aut}_{\text{aff}}(\Omega)$, so is S' . Choose $s_0 \in S$ and a sequence $\{g_n\} \subset \text{Aut}_{\text{aff}}(\Omega)$ converging to $g \in \overline{\text{Aut}(\Omega)} < \text{PM}(n+1, \mathbb{R}^n)$ such that $R(g) = s_0$. Then $K(g) = \partial\mathbb{R}^n$ and any point of $\mathbb{R}^n$ converges to s_0 . But all point of S' cannot converge to s_0 since $g_n(S') = S'$ for all g_n and S is an open subset of $\overline{S}$. Therefore we conclude $e \in S$. $\square$

5. STRICTLY CONVEX QUASI-HOMOGENEOUS DOMAINS

We will show in this section that there is only one strictly convex quasi-homogeneous affine domain up to affine equivalence, although there are infinitely many quasi-homogeneous strictly convex domains in $\mathbb{RP}^n$ as mentioned in the Section 2. To do this, we first investigate strictly convex quasi-homogeneous projective domains since every strictly convex quasi-homogeneous affine domain is a strictly convex quasi-homogeneous projective domain, as will be shown later in this section. For example, a strictly convex affine domain $D = \{(x, y) \in \mathbb{R}^2 \mid y > 1/x\}$ is not strictly convex when it is considered as a projective domain, and it is easy to show that D is not quasi-homogeneous.

5.1. Strictly convex quasi-homogeneous projective domains with twice differentiable boundary.

The aim of this section is to prove

Theorem 5.1. *Let Ω be a strictly convex quasi-homogeneous projective domain with twice differentiable boundary. Then Ω is projectively equivalent to a ball.*

This was already proved in two dimensional case by Kuiper [29] and Benzécri [7]. For arbitrary dimensional cases, we need some lemmas.

Lemma 5.2. *Let f and g be twice differentiable real-valued functions defined on $(-r, r)$. Suppose $f(0) = g(0)$, $f'(0) = g'(0)$ and $f(x) \geq g(x)$ for all $x \in (-r, r)$. Then $f''(0) \geq g''(0)$.*

Proof. If $f'(x) \leq g'(x)$ on $[0, a)$ for some $a > 0$, then $f - g$ is a non-increasing function on $[0, a)$. Since $f - g \geq 0$ and $(f - g)(0) = 0$ by hypothesis, $(f - g)(x) = 0$ on $[0, a)$ and this implies $f''(0) = g''(0)$.

Otherwise, for each integer $n > 0$ there exists a positive real number p_n such that $p_n < 1/n$ and $f'(p_n) > g'(p_n)$. Twice differentiability of $f - g$ implies the following equalities.

$$f''(0) - g''(0) = \lim_{h \rightarrow 0} \frac{f'(h) - g'(h)}{h} = \lim_{n \rightarrow \infty} \frac{f'(p_n) - g'(p_n)}{p_n}$$

Since for any n

$$\frac{f'(p_n) - g'(p_n)}{p_n} > 0,$$

we conclude $f''(0) - g''(0) \geq 0$. $\square$

Lemma 5.3. *Let f be a strictly convex twice differentiable function defined on an open subset D of $\mathbb{R}^n$. Then there exists a point $x \in D$ such that the Hessian of f at x is positive definite.*^a

Proof. We can choose a hemisphere which lies below and touch the graph of f at some $x \in D$. Let g be the defining function of the hemisphere. Then the Hessian of g is positive definite everywhere. By the above lemma, we see $D_v^2 f(x) \geq D_v^2 g(x) > 0$ for any direction v and this yields the desired result. $\square$

The proof of Theorem 5.1: First we choose an affine chart containing Ω . Let f be a local boundary equation of Ω . By Lemma 5.3, there exists a point p in $\partial\Omega$ such that the Hessian of f at p is positive definite. By Corollary 6 of Rockafellar

^aThe author learned the idea of the proof from Prof. Mohammad Ghomi.

[35], any twice differentiable convex function has a second-order Taylor expansion and thus we can take a suitable basis of $\mathbb{R}^n$ such that $p = (0, \dots, 0)$ and

$$\lim_{(x_1, \dots, x_{n-1}) \rightarrow 0} \frac{f(x_1, \dots, x_{n-1})}{x_1^2 + \dots + x_{n-1}^2} = 1.$$

This means that Ω has an osculating ellipsoid and so Ω is projectively equivalent to a ball by Theorems 2.7 and 2.8 (ii). $\square$

The above proof says in fact the following.

Proposition 5.4. *Let Ω be a strictly convex quasi-homogeneous projective domain whose boundary is twice differentiable on some open subset. Then Ω is projectively equivalent to a ball.*

5.2. Strictly convex quasi-homogeneous affine domains. In this section we will show that an n -dimensional paraboloid is the only strictly convex affine domain in $\mathbb{R}^n$ up to affine equivalence, as stated in the beginning of this section 5. We will obtain this result through the following steps.

- (I) Every strictly convex quasi-homogeneous affine domain has an 1-dimensional asymptotic cone, that is, it has an invariant direction.
- (II) The boundary of any strictly convex quasi-homogeneous affine domain is homogeneous and thus of class C^∞ .
- (III) A strictly convex quasi-homogeneous affine domain is affinely equivalent to a paraboloid.

Proposition 5.5. *Let Ω be a quasi-homogeneous properly convex affine domain in $\mathbb{R}^n$. Then Ω is strictly convex if and only if $AC(\Omega)$ is one-dimensional.*

Proof. Suppose Ω is strictly convex and $\dim AC(\Omega) \geq 2$. Consider Ω as a projective domain. Then $\overline{AC(\Omega)} \cap \mathbb{RP}_\infty^{n-1}$ is a closed face of Ω with $\dim \geq 1$, and thus $\Omega \subset \mathbb{R}^n$ has a line segment ℓ in $\partial_p \Omega$. (Note that this implies that Ω is not strictly convex as a projective domain.) From Theorem 2.8 (iv), we see Ω has a triangular section, and thus $\partial_a \Omega \cap \mathbb{R}^n$ has also a line segment. But this contradicts the fact that Ω is a strictly convex affine domain.




FIGURE 2.

To prove the other direction, suppose that $\dim(AC(\Omega)) = 1$ and Ω is not strictly convex. Suppose that there exists an open neighborhood $U \subset \mathbb{R}^n$ of $p \in \partial_a \Omega$ such that $\partial_a \Omega \cap U$ has no line segment. Then p is an extreme point of Ω and there exist $\{g_i\} \subset \text{Aut}(\Omega)$ such that $\lim_{i \rightarrow \infty} g_i = g$ and $R(g) = \{p\}$ by Lemma 3.4. Note that $\overline{\Omega} \cap \partial \mathbb{R}^n$ is one point set which equals to $\overline{AC(\Omega)} \cap \partial \mathbb{R}^n$. For any open subset V of $\mathbb{R}^n$, $g_i(V)$ converges uniformly to p , since $\mathbb{R}^n$ does not intersect $K(g)$ by Lemma 3.5. (In fact, $K(g)$ is exactly $\partial \mathbb{R}^n = \mathbb{RP}_\infty^{n-1}$.) Thus $g_i(V) \subset U$ for sufficiently large

i and this implies $g_i(V) \cap \partial_a \Omega$ has no line segment and so does $V \cap \partial_a \Omega$ if it is not empty. Since V is arbitrary we conclude that $\partial_a \Omega$ has no line segment, which is a contradiction. So Ω can not be strictly convex anywhere, i.e., any affine boundary point of Ω does not have a neighborhood containing no line segment.

Let L be an one dimensional linear subspace generated by $\text{AC}(\Omega)$. For each $x \in \Omega$, we defined $s(x)$ in section 4 by the point in $\partial_a \Omega$ satisfying the following:

$$s(x) + \text{AC}(\Omega) = \Omega \cap (x + L).$$

Recall that the map $s : \Omega \rightarrow \partial_a \Omega$ is continuous. Now choose an open set U contained in Ω . Then $s(U)$ is an open subset of $\partial_a \Omega$. Since Ω is not strictly convex everywhere, $s(U)$ contains an open line segment l . But by Lemma 3.6, $\partial_a \Omega$ can't have any nonzero dimensional bounded face. Therefore the asymptotic half line must be contained in the face containing l . But this contradicts that every asymptotic line starting at a point in $s(U)$ passes $U \subset \Omega$ (see the figure 2). $\square$

Remark 5.6. *This proposition implies that every strictly convex quasi-homogeneous affine domain is also strictly convex as a projective domain.*

Proposition 5.7. *Let Ω be a quasi-homogeneous affine domain in $\mathbb{R}^n$. If Ω is strictly convex, then its boundary is homogeneous.*

Proof. Let G be a subgroup of $\text{Aff}(n, \mathbb{R})$ which acts syndetically on Ω . By Proposition 5.5, there exists a one-dimensional vector space L such that $\text{AC}(\Omega) = \Omega \cap L$ is a half line. For each $x \in \Omega$, the intersection $\Omega_x = \Omega \cap (x + L)$ is a half line and $s(x)$ is the starting point of Ω_x . As stated in section 4, the map $s : \Omega \rightarrow \partial_a \Omega$ is continuous and induces a one-parameter group of homeomorphism of Ω by the following formula :

$$c_t(x) = s(x) + e^t(x - s(x)) \text{ for } t \in \mathbb{R}, x \in \Omega.$$

It is easy to show that c_t is d_H -decreasing for $t \geq 0$ (see figure 3) when Ω is strictly convex, that is,

$$d_H(c_t(x), c_t(y)) \leq d_H(x, y) \text{ for all } t \geq 0.$$




FIGURE 3.

On the other hand, we see that $g(\Omega_x) = \Omega_{gx}$ since L is an invariant direction, that is, since L is invariant under the action of the linear parts of $\text{Aut}_{\text{aff}}(\Omega)$. This implies

$$(2) \quad s(gx) = gs(x) \text{ for all } g \in G.$$

We may assume G is closed in $\text{Aff}(n, \mathbb{R})$. Then G acts on Ω properly by Proposition 3.7. Now we can define c_t on the orbit space $G \backslash \Omega$ by (2). In fact, for all $g \in G$

and $y \in \Omega$

$$\begin{aligned} c_t(gy) &= s(gy) + e^t(gy - s(gy)) = gs(y) + e^t(gy - gs(y)) \\ &= e^tgy + (1 - e^t)gs(y) = g(e^ty + (1 - e^t)s(y)) = gc_t(y) \end{aligned}$$

since every affine transformation preserves a convex combination. Note that Lemma 3.10 implies that there exists $g_{x,y} \in G$ such that

$$(3) \quad d_G(Gx, Gy) = d_H(x, g_{x,y}y)$$

where d_G is defined in (1).

It is obvious that for $t \geq 0$, the homeomorphisms c_t of $G \setminus \Omega$ are distance decreasing. By the way, the fact that G acts on Ω syndetically implies that $G \setminus \Omega$ is a compact metric space and it is well known that every distance decreasing surjection from a compact metric space onto itself is an isometry. But c_t is strictly distance decreasing on Ω for any pair (x, y) such that $s(x) \neq s(y)$, that is, $d_H(x, y) > d_H(c_t(x), c_t(y))$.

Suppose there exists a pair (x, y) such that $s(x) \neq gs(y)$ for all $g \in G$. Then we have

$$d_G(Gx, Gy) = d_H(x, g_{x,y}y) > d_H(c_t(x), c_t(g_{x,y}y)) = d_H(c_t(x), g_{x,y}c_t(y))$$

and this implies

$$d_G(Gx, Gy) > d_G(c_t(Gx), c_t(Gy)).$$

This is a contradiction. Therefore, for any pair (x, y) we get $s(x) = gs(y)$ for some $g \in G$ and so we conclude that $\partial_a \Omega$ is homogeneous. $\square$

Remark 5.8. We showed in the proof of the above proposition that the set of all asymptotic cone points S is homogeneous when it is strictly convex. In this case, $S = \partial \Omega$ since Ω has a 1-dimensional asymptotic cone. In case $\dim \text{AC}(\Omega)$ is not 1-dimensional, S is a proper subset of $\partial \Omega$. For example, if $\Omega = \{(x, y, z) \in \mathbb{R}^3 \mid x > 0, y > z^2\}$ then $S = \{(0, y, z) \in \mathbb{R}^3 \mid y = z^2\}$. S seems to be still homogeneous even in the case of $\dim \text{AC}(\Omega) > 1$, which implies that the Lie group $\text{Aut}(\Omega)$ is at least of dimension $n - \dim \text{AC}(\Omega)$. But this is not proved yet.

Now, we get the following main theorem of this section.

Theorem 5.9. Every strictly convex quasi-homogeneous affine domain in $\mathbb{R}^n$ is affinely equivalent to an n -dimensional paraboloid.

Proof. We have shown in Theorem 5.7 that $\partial_a \Omega$ is homogeneous and we can see $\partial_a \Omega$ is twice differentiable almost everywhere by Corollary 6 of [35] (in fact, $\partial_a \Omega$ is C^∞ since it is the orbit of the Lie group $\text{Aut}(\Omega)$). But the homogeneity of $\partial_a \Omega$ implies that $\partial_a \Omega$ is twice differentiable everywhere and thus Ω is projectively equivalent to a ball in $\mathbb{RP}^n$ by Proposition 5.4 since Ω is also strictly convex when it is considered as a projective domain. Therefore we conclude that Ω is affinely equivalent to a paraboloid since $\overline{\Omega} \cap \partial \mathbb{R}^n = \overline{\text{AC}(\Omega)} \cap \partial \mathbb{R}^n$ is a point by Proposition 5.5. $\square$

5.3. Strictly convex quasi-homogeneous projective domains. Using the results of the previous section, we will show that if Ω is a strictly convex quasi-homogeneous projective domain, then

- (i) $\partial \Omega$ is at least C^1 ,
- (ii) Ω is an ellipsoid if $\partial \Omega$ is twice differentiable,

(iii) if $\partial\Omega$ is C^α on an open subset of $\partial\Omega$, then $\partial\Omega$ is C^α everywhere.

Note that we already proved (ii) in Theorem 5.1. Now we prove (i).

Proposition 5.10. *Let Ω be a quasi-homogeneous properly convex domain in $\mathbb{RP}^n$. Then Ω is strictly convex if and only if the boundary of Ω is continuously differentiable.*

Proof. Note that for a convex function, the differentiability implies the continuity of the derivative (see Corollary 25.5.1 of [34]) and the existence of all partial derivatives implies the differentiability (see [37] p.27). So, to prove that $\partial\Omega$ is C^1 it suffices to show that $\partial\Omega$ has partial derivatives everywhere. Suppose that Ω is strictly convex and there exists a point $x \in \partial\Omega$ at which $\partial\Omega$ does not have some partial derivative. Then we can choose a 2-dimensional subspace H such that $H \cap \Omega$ is a 2-dimensional section of Ω containing x in its boundary and $\partial(H \cap \Omega)$ is not differentiable at x . This means that x is a conic face of $H \cap \Omega$, hence Ω has a triangular section by Theorem 2.8 (iv). But this contradicts the strict convexity of Ω .

Now we prove the converse. Suppose that $\partial\Omega$ is differentiable and Ω is not strictly convex. Since Ω is not strict convex, Ω has a triangular section by Corollary 2.9. This implies that $\partial\Omega$ can not be differentiable at any vertex point of the triangle. This contradiction completes the proof. $\square$

Proposition 5.11. *Let Ω be a quasi-homogeneous strictly convex domain in $\mathbb{RP}^n$. If $\partial\Omega$ is C^α on some neighborhood V of $p \in \partial\Omega$, then $\partial\Omega$ is C^α .*

Proof. We showed in Lemma 3.4 that for each point $p \in \partial\Omega$, there exists a sequence $\{g_n\} \subset \text{Aut}(\Omega)$ and $x \in \Omega$ such that $g_n x$ converges to p .

Obviously g is singular. By Lemma 3.2, $K(g) \cap \Omega = \emptyset$ and $R(g) \cap \Omega = \emptyset$. Strict convexity of Ω implies that the rank of g is 1 and $R(g) = \{p\}$ and $K(g) \cap \partial\Omega$ must be a single point, say q , by Lemma 3.2 again. For any $y \in \partial\Omega \setminus \{q\}$, $g(y) = p$ and thus $g_i(y) \in V$. Then $g_i^{-1}(V)$ is a C^α -neighborhood of y . Therefore $\partial\Omega$ is C^α except $q = K(g) \cap \overline{\Omega}$. Suppose that $\partial\Omega$ is not C^α at q . Then q must be a fixed point. Since $\partial\Omega$ is always C^1 by Proposition 5.10, there exists a unique supporting hyperplane H and thus H is invariant under $\text{Aut}(\Omega)$. So Ω must be a strictly convex quasi-homogeneous affine domain and so it is projectively equivalent to a ball by Theorem 5.9. This contradicts that $\partial\Omega$ is not C^α at q . Therefore $\partial\Omega$ is C^α everywhere. $\square$

From Propositions 5.10 and 5.11 and Theorem 5.1, we get immediately the following proposition.

Proposition 5.12. *Let Ω be a strictly convex quasi-homogeneous domain in $\mathbb{RP}^n$. Then either (i) Ω is an ellipsoid, or (ii) $\partial\Omega$ is C^1 and fails to be twice differentiable on a dense subset.*

A compact convex projective manifold M is a quotient of a convex projective domain Ω in $\mathbb{RP}^n$ by a discrete subgroup Γ of $\text{PGL}(n+1, \mathbb{R})$ acting on Ω properly discontinuously and freely. From the above proposition we get

Corollary 5.13. *If a strictly convex projective domain Ω covers a compact projectively flat manifold, then either Ω is an ellipsoid or $\partial\Omega$ fails to be twice differentiable on a dense subset.*

Remark 5.14. *This corollary was proved in the 2-dimensional case by Kuiper [29].*

The following proposition shows that there is no quasi-homogeneous strictly convex projective domain which is not divisible. In the 2-dimensional case, there is a more elementary and direct proof, which will be given in the next section. In fact, we will show later that every 3-dimensional quasi-homogeneous properly convex affine cone is divisible. It seems to be true that any irreducible properly convex projective domain is either homogeneous or divisible. Y. Benoist [5] proved this when the domain has an irreducible automorphism group.

Proposition 5.15. *Let Ω be a strictly convex quasi-homogeneous projective domain in $\mathbb{RP}^n$. Suppose that Ω is not an ellipsoid. Then $\text{Aut}(\Omega)$ is discrete and so Ω is a divisible domain.*

Proof. By the Proposition 4.2 of [5], it suffice to show that $\text{Aut}(\Omega)$ is irreducible. Suppose $\text{Aut}(\Omega)$ is reducible and L is a projective subspace of $\mathbb{RP}^n$ which is invariant under the action of $\text{Aut}(\Omega)$. Then $L \cap \Omega = \emptyset$ and $L \cap \overline{\Omega} \neq \emptyset$, since Ω is quasi-homogeneous. So the strict convexity of Ω implies that $L \cap \Omega$ must be a point $\{p\}$ in $\partial\Omega$. This implies p is a fixed point of $\text{Aut}(\Omega)$. Since every strictly convex quasi-homogeneous projective domain is of class C^1 , p has a unique tangent plane H_p and it is preserved by $\text{Aut}(\Omega)$. This means that Ω is a quasi-homogeneous strictly convex affine domain and so it must be an ellipsoid by Theorem 5.12, which gives a contradiction. $\square$

This Proposition shows that every strictly convex quasi-homogeneous domain covers a compact projectively flat manifold. In addition, W. Goldman and S. Choi [12] proved that any strictly convex $\mathbb{RP}^2$ -structure on a closed surface is a projective deformation of a hyperbolic structure. A 3-dimensional case was shown by I. Kim [24] and then Y. Benoist [6] has proved it for more general case.

6. QUASI-HOMOGENEOUS CONVEX AFFINE DOMAINS OF DIMENSION ≤ 3

In this section, we will characterize low dimensional quasi-homogeneous convex affine domains which are not cones.

Proposition 6.1. *Let Ω be a properly convex quasi-homogeneous affine domain in $\mathbb{R}^2$. Then Ω is affinely equivalent to either a quadrant or a parabola.*

Proof. If $\dim \text{AC}(\Omega) = 1$ then Ω is affinely equivalent to a parabola by Proposition 5.5 and Theorem 5.9. Otherwise $\dim \text{AC}(\Omega) = 2$, and hence Ω must be a properly convex cone by Corollary 4.5. Any two dimensional properly convex cone is affinely equivalent to a quadrant, which completes the proof. $\square$

Proposition 6.2. *Let Ω be a properly convex quasi-homogeneous affine domain in $\mathbb{R}^3$. Suppose Ω is not a cone. Then Ω is affinely equivalent to one of the following:*

- (i) *A 3-dimensional paraboloid, i.e., $\{(x, y, z) \in \mathbb{R}^3 \mid z > x^2 + y^2\}$.*
- (ii) *A convex domain bounded by a parabola $\times \mathbb{R}^+$, i.e., $\{(x, y) \in \mathbb{R}^2 \mid y > x^2\} \times \{z \in \mathbb{R} \mid z > 0\} = \{(x, y, z) \in \mathbb{R}^3 \mid y > x^2, z > 0\}$.*

Proof. By Theorem 5.9 and Proposition 5.5, Ω is affinely equivalent to a paraboloid if $\dim \text{AC}(\Omega) = 1$. Since Ω is not a cone, $\dim \text{AC}(\Omega) \neq 3$ by Corollary 4.5. Hence it suffices to show that Ω is affinely equivalent to a convex domain bounded by a parabola $\times \mathbb{R}^+$ if $\dim \text{AC}(\Omega) = 2$. We know that Ω admits a parallel foliation by cosets of the asymptotic cone of Ω by Proposition 4.4. Put $\text{AC}_\infty(\Omega) = \overline{\text{AC}(\Omega)} \cap \mathbb{RP}_\infty^2 = \overline{ab}$. Now we will show that one of $\{a, b\}$ is a conic

point of Ω . Choose an asymptotic cone point $p \in \partial\Omega$, then the triangle $\triangle pab$ is a 2-dimensional section of Ω . Therefore there exists supporting hyperplanes H_{pa} and H_{pb} tangent to $\overline{pa}$ and $\overline{pb}$ respectively. Let $L = H_{pa} \cap H_{pb}$. Suppose every asymptotic cone point is contained in L . Then the fact that Ω is not a cone implies that the set of all asymptotic cone points in $L \cap \partial\Omega$ cannot be one point set and thus it is a bounded line, since L is not an asymptotic direction. This contradicts Lemma 3.6. So, there exists an asymptotic cone point $p' \in \partial\Omega$ such that $p' \notin L$. This means that $p' \notin H_{pa}$ or $p' \notin H_{pb}$. We may assume that $p' \notin H_{pa}$. Note that H_{pa} and $H_{p'a}$ are neither the same hyperplane nor parallel. So a is a conic face of Ω when it is considered as a projective domain. In fact a is the intersection of the three independent supporting hyperplanes $H_{pa}, H_{p'a}, H_{ab}$, where H_{ab} is in fact $\partial\mathbb{R}^3 = \mathbb{RP}_\infty^2$. By Theorems 2.7 and 2.8 (i), we conclude that $\Omega = a + \Omega'$ for some 2-dimensional quasi-homogeneous properly convex projective domain Ω' . Ω' must contain all asymptotic cone points of Ω and be an invariant quasi-homogeneous 2-dimensional affine domain with 1-dimensional asymptotic cone. By Theorem 6.1, Ω' is a parabola. Consequently, we have that $\Omega = \overline{pa} \times \Omega'$. $\square$

If Ω is a quasi-homogeneous convex affine domain which is not properly convex, then Ω is a product $\Omega' \times \mathbb{R}^k$ with $k > 0$, where Ω' is a $(n - k)$ -dimensional quasi-homogeneous properly convex affine domain. So we get the following two corollaries from Propositions 6.1 and 6.2.

Corollary 6.3. *Let Ω be a quasi-homogeneous convex affine domain in $\mathbb{R}^2$. Then Ω is affinely equivalent to one of the following :*

- (i) $\{(x, y) \in \mathbb{R}^2 \mid y > x^2\}$,
- (ii) $\{(x, y) \in \mathbb{R}^2 \mid x > 0, y > 0\}$,
- (iii) $\{(x, y) \in \mathbb{R}^2 \mid y > 0\}$,
- (iv) $\mathbb{R}^2$.

Every 3-dimensional quasi-homogeneous properly convex cone is either a strictly convex cone or a simplex cone since a 2-dimensional quasi-homogeneous properly convex projective domain which is not a triangle is strictly convex. So we get

Corollary 6.4. *Let Ω be a convex quasi-homogeneous domain in $\mathbb{R}^3$. The Ω is affinely equivalent to one of the following :*

- (i) $\{(x, y, z) \in \mathbb{R}^3 \mid z > x^2 + y^2\}$,
- (ii) $\{(x, y, z) \in \mathbb{R}^3 \mid y > x^2, z > 0\}$,
- (iii) $\{(x, y, z) \in \mathbb{R}^3 \mid x > 0, y > 0\}$,
- (iv) $\{(x, y, z) \in \mathbb{R}^3 \mid y > x^2\}$,
- (v) $\{(x, y, z) \in \mathbb{R}^3 \mid z > 0\}$,
- (vi) $\mathbb{R}^3$,
- (vii) a quasi-homogeneous strictly convex affine cone,
- (viii) a simplex cone.

Remark 6.5. (i) *Every 3-dimensional quasi-homogeneous properly convex affine cone is divisible. The reason is as follows: Any quasi-homogeneous properly convex 2-dimensional projective domain which is neither a triangle nor an ellipsoid is a strictly convex domain whose boundary is not twice differentiable on a dense subset by Proposition 5.12. If the automorphism group of a 2-dimensional strictly convex domain has a 1-dimensional Lie subgroup, the boundary has a 1-dimensional orbit and consequently is twice differentiable, that is, the boundary is twice differentiable on some open set. So the*

automorphism group of a 2-dimensional quasi-homogeneous strictly convex domain whose boundary is not twice differentiable, must be discrete. Therefore every 2-dimensional quasi-homogeneous properly convex projective domain is divisible and so is any 3-dimensional quasi-homogeneous properly convex affine cone. Actually, this fact follows from Proposition 5.15.

- (ii) *Any strictly convex cone of (vii) is either an elliptic cone $\{(x, y, z) \in \mathbb{R}^3 \mid x^2 - y^2 - z^2 > 0\}$ or its boundary is not twice differentiable on any open subset by Proposition 5.12.*

7. COMPACT CONVEX AFFINE MANIFOLDS OF DIMENSION ≤ 4

Let M be a compact convex affine manifold. Then we get a divisible domain Ω and a discrete subgroup Γ of $\text{Aut}_{\text{aff}}(\Omega)$ such that $\Omega/\Gamma = M$. If Ω is properly convex, then it must be a cone by a result of Vey [42].

For a 2-dimensional compact convex affine manifold M , we know that Ω is a quadrant if it is properly convex. Otherwise Ω is a half space since the only 1-dimensional quasi-homogeneous affine domain is $\mathbb{R}^+$.

Using the results in the previous section, we get the following.

Theorem 7.1. *Let Ω be an affine domain in $\mathbb{R}^3$ which covers a compact convex affine manifold. Then Ω is affinely equivalent to one of the following :*

- (i) $\mathbb{R}^3$,
- (ii) $\mathbb{R}^2 \times \mathbb{R}^+$,
- (iii) $\mathbb{R} \times \{(x, y) \in \mathbb{R}^2 \mid y > x^2\}$,
- (iv) $\mathbb{R} \times \{(x, y) \in \mathbb{R}^2 \mid x > 0, y > 0\}$,
- (v) *a quasi-homogeneous strictly convex cone,*
- (vi) *a simplex cone.*

Proof. Consider a 3-dimensional quasi-homogeneous convex affine domain Ω . It is isomorphic to $\mathbb{R}^k \times \Omega'$ for some $(3 - k)$ -dimensional properly convex affine domain Ω' . If $k = 3$, $\Omega = \mathbb{R}^3$ and if $k = 2$, $\Omega = \mathbb{R}^2 \times \mathbb{R}^+$. Since a 2-dimensional quasi-homogeneous properly convex domain is either a quadrant or a parabola, Ω is either (iii) or (iv) if $k = 1$. If $k = 0$, that is, if Ω is properly convex, then it is a cone and thus either (v) or (vi) by Corollary 6.4.

All of these quasi-homogeneous domains are actually divisible. The divisibility of (iii) was shown by W. Goldman in [20] and the divisibility of (v) is implied by Remark 6.5. The remaining cases are easy. $\square$

Similarly, we get the following theorem in 4-dimensional case.

Theorem 7.2. *Let Ω be an affine domain in $\mathbb{R}^4$ which covers a compact convex affine manifold. Then Ω is affinely equivalent to one of the following :*

- (i) $\mathbb{R}^4$,
- (ii) $\mathbb{R}^3 \times \mathbb{R}^+$,
- (iii) $\mathbb{R}^2 \times \{(x, y) \in \mathbb{R}^2 \mid y > x^2\}$,
- (iv) $\mathbb{R}^2 \times \{(x, y) \in \mathbb{R}^2 \mid x > 0, y > 0\}$,
- (v) $\mathbb{R} \times \{(x, y, z) \in \mathbb{R}^3 \mid z > x^2 + y^2\}$,
- (vi) $\mathbb{R} \times \mathbb{R}^+ \times \{(x, y) \in \mathbb{R}^2 \mid y > x^2\}$,
- (vii) $\mathbb{R} \times$ *a divisible properly convex cone of dimension 3,*
- (viii) *a divisible properly convex cone of dimension 4*

W. Goldman and M. Hirsch proved in [22] that any compact complete affine 3-manifold has parallel volume (see Cor. 3.3). They also proved that if a compact affine n -manifold M with holonomy group Γ has parallel volume, then the algebraic hull $A(\Gamma)$ of Γ acts transitively and so Γ preserves no proper algebraic subset of $\mathbb{R}^n$. Considering the developing image Ω of compact convex affine manifold of dimension ≤ 4 , we see that the holonomy group of compact convex affine manifold has a proper invariant algebraic set if its developing map is not surjective. So we get the following corollary.

Corollary 7.3. *Let M be a compact convex affine n -manifold.*

- (i) *For $n = 3$, M is complete if and only if it has parallel volume.*
- (ii) *For $n = 4$, M is complete if it has parallel volume.*

ACKNOWLEDGEMENTS

I would like to thank Prof. Hyuk Kim who has encouraged and advised me all the while I have been studying this topic. I am also grateful to Prof. Inkang Kim for reading this article and helpful discussions and Prof. Suhyoung Choi for his critical comments. Conversations with Prof. Yves Benoist and Mohammad Ghomi by e-mail have been also helpful. Finally I thank the referee and Prof. Seong-deog Yang for suggesting a number of improvements on the exposition.

REFERENCES

- [1] T. Barbot, *Variétés affines radiales de dimension trois*, Bull. Soc. Math. France **128** (2000), no 3, 347–389.
- [2] Y. Benoist, *Automorphismes des cônes convexes*, Inv. Math. **141** (1) (2000), 149–193.
- [3] Y. Benoist, *Convex divisibles*, Comp. Rend. Ac. Sc. **332** (5) (2001) p.387–390.
- [4] Y. Benoist, *Convex divisibles I*, preprint (2001).
- [5] Y. Benoist, *Convex divisibles II*, to appear in Duke Math. Journ.
- [6] Y. Benoist, *Convex divisibles III*, preprint (2001).
- [7] J. P. Benzécri, *Sur les variétés localement affines et projectives*, Bull. Soc. Math. France **88** (1960), 229–332.
- [8] Y. V. Carrière, *Un survol de la théorie des variétés affines*, Séminaire de théorie spectrale et géométrie, CHAMBERY-GRENOBLE, (1987–1988), 9–22.
- [9] Y. V. Carrière, *Autour de la conjecture de L. Markus sur les variétés affines*, Invent. Math. **95** (1989), 615–628.
- [10] Y. V. Carrière, *Autour de la conjecture de L. Markus sur les variétés affines*, Invent. Math. **95** (1989), 615–628.
- [11] H. Cho, *A classification of convex affine 3-manifold*, Ph.D thesis, Seoul National University, (1999).
- [12] S. Choi and W. M. Goldman, *Convex real projective structures on closed surfaces are closed*, Proc. Amer. Math. Soc. **118** (1993), 657–661.
- [13] B. Colbois and P. Verovic, *Rigidity of Hilbert metrics*, Bull. Austral. Math. Soc. **65** (2002), no.1, 23–34.
- [14] D. Fried, *Distality, completeness and affine structures*, J. Diff. Geom., **24** (1986), 265–273.
- [15] D. Fried, W. M. Goldman and M. Hirsch, *Affine manifolds with nilpotent holonomy*, Comment. Math. Helv. **56** (1981), 487–523.
- [16] D. Fried and W. Goldman, *Three-dimensional affine crystallographic groups*, Adv. in Math. **47** (1983), no.1, 1–49.
- [17] W. M. Goldman, *Geometric structures on manifolds and varieties of representations*, Contemp. Math. **74** (1988), 169–198.
- [18] W. M. Goldman, *Convex real projective structures on compact surfaces*, J. Diff. Geom. **31** (1990), 791–845.
- [19] W. M. Goldman, *Projective geometry on manifolds*, Lecture Notes (v.0.3)- Mathematics 748B Spring (1988).

- [20] W. M. Goldman, *Two examples of affine manifolds*, Pacific J. Math. **94** (1981), no.2, 327–330.
- [21] W. M. Goldman and M. W. Hirsch, *The radiance obstruction and parallel forms on affine manifolds*, Trans. Amer. Math. Soc. **286** (1984), 629–649.
- [22] W. M. Goldman and M. W. Hirsch, *Affine manifolds and orbits of algebraic groups*, Trans. Amer. Math. Soc. **295** (1986), no.1, 175–198.
- [23] H. Kim, *Actions of infinite discrete groups of projective transformations*, Proceedings of the second GARC Symposium on pure and applied mathematics, Part III, Lecture Note series (1993), no.17, 1–9.
- [24] I. Kim, *Rigidity and deformation spaces of strictly convex real projective structures on compact manifolds*, J. Diff. Geom. **58** (2) (2001), 189–218.
- [25] S. Kobayashi, *Intrinsic distances associated with flat affine or projective structures*, J. Fac. Sci. Univ. Tokyo **24** (1) (1977), 129–135.
- [26] S. Kobayashi, *Projectively invariant distances for affine and projective structures*, Differential Geometry, Banach Center Publ. **12** (1979), 127–152.
- [27] S. Kobayashi, *Invariant distances for projective structures*, Symposia Mathematica **XXVI**, Rome (1982), 153–161.
- [28] J. L. Koszul, *Déformation des connexions localement plats*, Ann. Inst. Fourier **18** (1968), 103–114.
- [29] N. H. Kuiper, *On convex locally projective spaces*, Convegno Intern. Geom. Diff. Italy (1953), 200–213.
- [30] L. Markus, *Cosmological models in differential geometry*, mimeographed notes, Univ. of Minnesota (1962), p.58.
- [31] T. Nagano and K. Yagi, *The affine structures on the real two-torus*, Osaka J. Math. **11** (1974), 181–210.
- [32] K. S. Park, *Invariant open sets under cocompact affine actions*, Bull. Korean Math. Soc. **36** (1) (1999), 203–207.
- [33] J. G. Ratcliffe, *Foundations of Hyperbolic Manifolds*, Springer-Verlag New York, Inc. (1994).
- [34] R. T. Rockafellar, *Convex analysis*, Princeton University Press (1970).
- [35] R. T. Rockafellar, *Second derivatives in convex analysis*, Nonlinear analysis and convex analysis: Proceedings of the international conference on Niigata, Japan, 28–31 July (1998), 3–7.
- [36] W. Rudin, *Functional analysis*, International series in Pure and Applied Mathematics (1991), p.75.
- [37] R. Schneider, *Convex bodies : the Brunn-Minkowski theory*, Cambridge University Press (1993).
- [38] J. Smillie, *Affinely flat manifolds*, Doctoral Dissertation, University of Chicago (1977).
- [39] W. P. Thurston, *The geometry and topology of 3-manifolds*, Preprint (1977).
- [40] J. Vey, *Une notion d'hyperbolicité sur les variétés localement plates*, C. R. Acad. Sci. Paris Ser. AB **266** (1968), A622–A624.
- [41] J. Vey, *Sur les automorphismes affines des ouverts convexes dans les espaces numériques*, C. R. Acad. Sci. Paris Ser. AB **270** (1970), A249–A251.
- [42] J. Vey, *Sur les automorphismes affines des ouverts convexes saillants*, Ann. Scuola Norm. Sup. Pisa **24** (3) (1970), 641–665.
- [43] E. B. Vinberg and V. G. Kats, *Quasi-homogeneous cones*, Math. Notes **1** (1967), 231–235. (translated from Math. Zametki **1** (1967), 347–354.)

APPLIED MATHEMATICS, SEJONG UNIVERSITY, 143-747, SEOUL, KOREA
 E-mail address: khjo@sejong.ac.kr